

### Written Assignment 3: Due Wednesday, September 28

**Problem 1:** Consider the linear transformation  $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$  given by:

$$T\left(\begin{pmatrix} x \\ y \end{pmatrix}\right) = \begin{pmatrix} x - y \\ x + y \end{pmatrix}.$$

Is  $T$  injective? Justify your answer carefully.

**Problem 2:** Let  $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a linear transformation. Recall that

$$\text{range}(T) = \{\vec{w} \in \mathbb{R}^2 : \text{There exists } \vec{v} \in \mathbb{R}^2 \text{ with } \vec{w} = T(\vec{v})\}.$$

Notice that  $\vec{0} \in \text{range}(T)$  because we know that  $T(\vec{0}) = \vec{0}$  by Proposition 2.4.2.

- Show that if  $\vec{w}_1, \vec{w}_2 \in \text{range}(T)$ , then  $\vec{w}_1 + \vec{w}_2 \in \text{range}(T)$ .
- Show that if  $\vec{w} \in \text{range}(T)$  and  $c \in \mathbb{R}$ , then  $c\vec{w} \in \text{range}(T)$ .

**Problem 3:** We defined linear transformations from  $\mathbb{R}^2$  to  $\mathbb{R}^2$ , but we can also define them from  $\mathbb{R}$  to  $\mathbb{R}$  as follows. A linear transformation from  $\mathbb{R}$  to  $\mathbb{R}$  is a function  $f: \mathbb{R} \rightarrow \mathbb{R}$  with both of the following properties:

- $f(x + y) = f(x) + f(y)$  for all  $x, y \in \mathbb{R}$ .
- $f(c \cdot x) = c \cdot f(x)$  for all  $c, x \in \mathbb{R}$ .

- Let  $r \in \mathbb{R}$ . Show that the function  $g_r: \mathbb{R} \rightarrow \mathbb{R}$  given by  $g_r(x) = rx$  is a linear transformation
- Show that if  $f: \mathbb{R} \rightarrow \mathbb{R}$  and  $g: \mathbb{R} \rightarrow \mathbb{R}$  are both linear transformations, and  $f(1) = g(1)$ , then  $f = g$ .