Problem Set 21: Due Friday, May 3

Problem 1: Let V be the vector space of all 2×2 matrices. Define $T: V \to V$ by letting

$$T\left(\begin{pmatrix} a & b \\ c & d \end{pmatrix}\right) = \begin{pmatrix} a & c \\ b & d \end{pmatrix}.$$

Notice that the function T takes an input matrix and outputs the result of switching the rows and columns (which is called the transpose of the original matrix). It turns out that T is a linear transformation. Let

$$\alpha = \left(\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \right),$$

and recall that α is a basis for V. What is $[T]^{\alpha}_{\alpha}$? Explain briefly.

Problem 2: Consider the unique linear transformation $T: \mathbb{R}^4 \to \mathbb{R}^3$ with

$$[T] = \begin{pmatrix} 1 & 0 & 2 & -1 \\ 3 & 1 & 9 & -5 \\ -1 & 2 & 4 & -2 \end{pmatrix}.$$

- a. Find (with explanation) bases for each of range(T) and Null(T).
- b. Calculate rank(T) and nullity(T).

Problem 3: Consider the unique linear transformation $T: \mathbb{R}^4 \to \mathbb{R}$ with

$$[T] = (0 \quad 1 \quad -3 \quad 7).$$

- a. Find (with explanation) bases for each of range(T) and Null(T).
- b. Calculate rank(T) and nullity(T).

Problem 4: Define $T: \mathcal{P}_5 \to \mathcal{P}_5$ by letting T(f) = f'', i.e. T(f) is the second derivative of f. Determine, with explanation, both rank(T) and nullity(T).

Problem 5: Let V be the vector space of all 2×2 matrices. Explain why there is no surjective linear transformation $T: V \to \mathcal{P}_4$.