

Homework 3: Due Wednesday, February 15

Problem 1: Consider the set $\mathbb{R}^{\geq 0} = \{x \in \mathbb{R} : x \geq 0\}$ of nonnegative reals. Let $*$ be the binary operation on $\mathbb{R}^{\geq 0}$ given by exponentiation, i.e. $a * b = a^b$.

- Is $*$ an associative operation on $\mathbb{R}^{\geq 0}$? Either prove or give an explicit counterexample.
- Does $*$ have an identity on $\mathbb{R}^{\geq 0}$? Prove your answer.

Problem 2: Define a binary operation $*$ on \mathbb{R} by letting $a * b = a + b + ab$.

- Show that $*$ is commutative, i.e. that $a * b = b * a$ for all $a, b \in \mathbb{R}$.
- Show that $*$ is associative, i.e. that $(a * b) * c = a * (b * c)$ for all $a, b, c \in \mathbb{R}$.
- Show that \mathbb{R} with operation $*$ has an identity element.
- Show that the set of invertible elements of $*$ equals $\mathbb{R} \setminus \{-1\} = \{x \in \mathbb{R} : x \neq -1\}$.

Note: Using Corollary 4.3.5, it follows that $\mathbb{R} \setminus \{-1\}$ under $*$ and with the identity element from part (c) is an abelian group.

Problem 3: Let S be the set of all 2×2 matrices of the form

$$\begin{pmatrix} a & a \\ a & a \end{pmatrix}$$

where $a \in \mathbb{R}$ and $a \neq 0$.

- Show that if $A, B \in S$, then $AB \in S$. Thus, matrix multiplication is a binary operation on S .
- Show that S with matrix multiplication has an identity element.
- Notice that every matrix in S has determinant 0, so every matrix in S fails to be invertible in the linear algebra sense. Nevertheless, show that S is a group under matrix multiplication with the identity from part (b).

Problem 4: Use the Euclidean Algorithm to show that $\overline{153} \in U(\mathbb{Z}/385\mathbb{Z})$ and to explicitly find its inverse.

Problem 5:

- Write out the Cayley table of $U(\mathbb{Z}/18\mathbb{Z})$.
- Compute (with explanation) the order of $\overline{11}$ in $U(\mathbb{Z}/18\mathbb{Z})$.

Problem 6: Let G be a group. Show that G is abelian if and only if $(a \cdot b)^{-1} = a^{-1} \cdot b^{-1}$ for all $a, b \in G$.

Problem 7: Suppose that $n \geq 3$. Let $\sigma \in S_n$ with $\sigma \neq id$. Show that there exists $\tau \in S_n$ such that $\sigma\tau \neq \tau\sigma$.

Hint: I strongly recommend that you avoid cycle notation and just work with σ as a function. Since $\sigma \neq id$, start by fixing an i with $\sigma(i) \neq i$. Now build a function $\tau \in S_n$ and show that it works.